arXiv:1512.01754v2 [math.GN] 1 Jun 2016

ON STABLE BAIRE CLASSES

OLENA KARLOVA AND VOLODYMYR MYKHAYLYUK

Abstract. We introduce and study adhesive spaces. Using this concept we obtain a characterization of stable Baire maps $f : X \to Y$ of the class α for wide classes of topological spaces. In particular, we prove that for a topological space X and a contractible space Y a map $f : X \to Y$ belongs to the n'th stable Baire class if and only if there exist a sequence $(f_k)_{k=1}^{\infty}$ of continuous maps $f_k : X \to Y$ and a sequence $(F_k)_{k=1}^{\infty}$ of functionally ambiguous sets of the n'th class in X such that $f|_{F_k} = f_k|_{F_k}$ for every k. Moreover, we show that every monotone function $f : \mathbb{R} \to \mathbb{R}$ is of the α'th stable Baire class if and only if it belongs to the first stable Baire class.

1. Introduction, terminology and notations

We say that a sequence $(f_n)_{n=1}^{\infty}$ of maps $f_n : X \to Y$ between topological spaces X and Y is *stably convergent to a map* $f : X \to Y$ *on* X, if for every $x \in X$ there exists $k \in \mathbb{N}$ such that $f_n(x) = f(x)$ for all $n \geq k$. A map $f : X \to Y$ belongs to *the first stable Baire class*, if there exists a sequence of continuous maps between X and Y which is stably convergent to f on X.

Real-valued functions of the first stable Baire class on a topological space X naturally appear both as an interesting subclass of all differences of semi-continuous functions on X [6, 7] and in problems on the Baire classification of integrals depending on a parameter [2] as well as in problems concerning a composition of Baire functions [8].

Real-valued functions of higher stable Baire classes were introduced and studied by Á. Császár and M. Laczkovich in [4, 5]. A characterization of maps of the first Baire class defined on a perfectly paracompact hereditarily Baire space with the Preiss-Simon property and with values in a path-connected space with special extension properties was established by T. Banakh and B. Bokalo in [1].

This paper is devoted to obtain a characterization of stable Baire maps for wide classes of topological spaces and any ordinal $\alpha \in [1, \omega_1)$. To do this we introduce a class of adhesive spaces and study their properties in Section 2. In Section 3 we give a characterization of stable Baire maps defined on a topological space and with values in adhesive spaces (see Theorem 3.2). Finally, in Section 4 we apply this characterization to classify monotone functions within stable Baire classes (see Theorem 4.2).

Let us give some notations and recall several definitions. For topological spaces X and Y by $\mathrm{C}(X, Y)$ we denote the collection of all continuous maps between X and Y.

1991 *Mathematics Subject Classification.* Primary 54C08, 54H05; Secondary 26A21.
Key words and phrases. stable convergence, stable Baire class, adhesive space.

If $A \subseteq Y^X$, then the symbol $\overline{A}^{\,\mathrm{st}}$ stands for the collection of all stable limits of sequences of maps from A. We put

$$\mathrm{B}_0^{\mathrm{st}}(X,Y) = \mathrm{C}(X,Y)$$

and for each ordinal $\alpha \in (0,\omega_1)$ let $\mathrm{B}_\alpha^{\mathrm{st}}(X,Y)$ be the family of all maps of *the α'th stable Baire class* which is defined by the formula

$$\mathrm{B}_\alpha^{\mathrm{st}}(X,Y) = \overline{\bigcup_{\beta<\alpha} \mathrm{B}_\beta^{\mathrm{st}}(X,Y)}^{\,\mathrm{st}}.$$

Recall that a set $A \subseteq X$ is *functionally closed*, if there exists a continuous function $f : X \to [0,1]$ with $A = f^{-1}(0)$. If the complement of A is functionally closed, then A is called *functionally open*.

Let $\mathcal{M}_0(X)$ be the family of all functionally closed subsets of X and let $\mathcal{A}_0(X)$ be the family of all functionally open subsets of X. For every $\alpha \in [1,\omega_1)$ we put

$$\mathcal{M}_\alpha(X) = \Big\{ \bigcap_{n=1}^{\infty} A_n : A_n \in \bigcup_{\beta<\alpha} \mathcal{A}_\beta(X),\ n = 1,2,\dots \Big\} \text{ and}$$

$$\mathcal{A}_\alpha(X) = \Big\{ \bigcup_{n=1}^{\infty} A_n : A_n \in \bigcup_{\beta<\alpha} \mathcal{M}_\beta(X),\ n = 1,2,\dots \Big\}.$$

Elements from the class $\mathcal{M}_\alpha(X)$ belong to *the α'th functionally multiplicative class* and elements from $\mathcal{A}_\alpha(X)$ belong to *the α'th functionally additive class* in X. We say that a set is *functionally ambiguous of the α'th class*, if it belongs to $\mathcal{M}_\alpha(X)$ and $\mathcal{A}_\alpha(X)$ simultaneously.

A topological space X is called *contractible*, if there exist a point $x_0 \in X$ and a continuous map $\gamma : X \times [0,1] \to X$ such that $\gamma(x,0) = x$ and $\gamma(x,1) = x_0$ for all $x \in X$. A space Y is *an extensor for X*, if for any closed set $F \subseteq X$ each continuous map $f : F \to Y$ can be extended to a continuous map $g : X \to Y$.

A map $f : X \to Y$ is called *piecewise continuous*, if there exists a cover $(F_n : n \in \mathbb{N})$ of X by closed sets such that each restriction $f|_{F_n}$ is continuous.

We denote by symbols $C(f)$ and $D(f)$ the sets of all points of continuity and discontinuity of a map $f : X \to Y$, respectively.

2. Adhesive spaces

Definition 2.1. We say that a topological space Y is *an adhesive for X* (we denote this fact by $Y \in \mathrm{Ad}(X)$), if for any two disjoint functionally closed sets A and B in X and any continuous maps $f, g : X \to Y$ there exists a continuous map $h : X \to Y$ such that $h|_A = f|_A$ and $h|_B = g|_B$. A space Y is said to be *an absolute adhesive* for a class $\mathcal{C}$ of topological spaces and is denoted by $Y \in \mathrm{AAd}(\mathcal{C})$, if $Y \in \mathrm{Ad}(X)$ for any $X \in \mathcal{C}$.

If the case when Y is an adhesive for any space X, then we will write $Y \in \mathrm{AAd}$.

Remark 2.2. (1) Every extensor is an adhesive.

(2) Let $\pi(X,Y)$ be the set of all homotopy classes of continuous maps between X and Y. If $\pi(X,Y) = \mathrm{C}(X,Y)$, then Y is an adhesive for X.

(3) Example 2.9(a) shows that the class of adhesive spaces is strictly wider than the class of extensors. Example 2.9(b) contains an adhesive Y for X such that $\pi(X,Y) \neq \mathrm{C}(X,Y)$.

Definition 2.3. Let E be a subspace of a topological space Y. The pair (Y, E) is called *a $*$-adhesive for X* (and is denoted by $(Y, E) \in \mathrm{Ad}^*(X)$), if there exists a point $y^* \in E$ such that for any continuous map $f : X \to Y$ and any two disjoint functionally closed sets A and B in X with $f(A) \subseteq E$ there exists a continuous map $h : X \to Y$ such that $h|_A = f|_A$ and $h|_B = y^*$.

Remark 2.4. A pair (Y, E) is a $*$-adhesive for X, if

(1) E is a subspace of $Y \in \mathrm{Ad}(X)$;
(2) $(Y, E) \in \mathrm{AE}(X)$ (i.e., each continuous map $f : F \to E$ has a continuous extension $g : X \to Y$);
(3) $E \in \mathrm{Ad}(X)$ and E is a retract of Y.

Definition 2.5. A topological space Y is said to be *a σ-adhesive for X*, if there exists a cover $(Y_n : n \in \mathbb{N})$ of Y by functionally closed subspaces Y_n such that $(Y, Y_n) \in \mathrm{Ad}^*(X)$ for every n.

Definition 2.6. A topological space X is *low-dimensional*, if each point of X has a base of open neighborhoods with discrete (probably, empty) boundaries.

It is clear that for a regular low-dimensional space we have $\mathrm{ind}X \leq 1$.

The following result gives examples of adhesive spaces.

Theorem 2.7. *Let Y be a topological space. Then*

(1) *Y is an absolute adhesive for the class of all strongly zero-dimensional spaces;*
(2) *Y is an absolute adhesive for the class of all low-dimensional compact Hausdorff spaces, if Y is path-connected;*
(3) *$Y \in \mathrm{AAd}$ if and only if Y is contractible.*

Proof. Let X be a topological space, A, B be disjoint functionally closed subsets of X and $f, g : X \to Y$ be continuous maps.

(1). Assume that X is strongly zero-dimensional and choose a clopen set $U \subseteq X$ such that $A \subseteq U \subseteq X \setminus B$. Then the map $h : X \to Y$,

$$h(x) = \begin{cases} f(x), & x \in U, \\ g(x), & x \in X \setminus U, \end{cases}$$

is continuous, $h|_A = f|_A$ and $h|_B = g|_B$.

(2). Assume that X is a low-dimensional Hausdorff compact space and Y is a path-connected space. We choose a continuous function $\varphi : X \to [0, 1]$ such that $A = \varphi^{-1}(0)$ and $B = \varphi^{-1}(1)$. For each point $x \in A \cup B$ we take its open neighborhood O_x with the discrete boundary ∂O_x such that $O_x \subseteq \varphi^{-1}([0, 1/3))$ for $x \in A$ and $O_x \subseteq \varphi^{-1}((2/3, 1])$ for $x \in B$. Since X is compact, every boundary ∂O_x is finite. Moreover, there exist finite subcovers $\mathcal{U}$ and $\mathcal{V}$ of $(O_x : x \in A)$ and $(O_x : x \in B)$, respectively. Then the sets $U = \cup\mathcal{U}$ and $V = \cup\mathcal{V}$ are open neighborhoods of A and B, respectively, $\partial U \cap \partial V = \emptyset$ and $|\partial U \cup \partial V| < \aleph_0$. In the case when one of the boundaries of the sets U or V is empty, we can construct the required map h as in case (1)). Hence, we may suppose that $\partial U \neq \emptyset \neq \partial V$. Let $\partial U = \{x_1, \ldots, x_n\}$ and $\partial V = \{x_{n+1}, \ldots, x_{n+m}\}$ for some $n, m \in \mathbb{N}$. Taking into account that the space $D = \partial U \cup \partial V$ is finite and Hausdorff, we obtain that a function $\psi : D \to [1, n + m]$, which is defined by the equality $\psi(x_i) = i$ for $i \in \{1, \ldots, n+m\}$, is continuous. Let $\tilde{\psi} : X \to [1, n+m]$ be a continuous extension of ψ. Denote $y_i = f(x_i)$ for $i \in \{1, \ldots, n\}$ and $y_i = g(x_i)$ for $i \in \{n+1, \ldots, n+m\}$.

Now we use the path-connectedness of Y and for every $i \in \{1, \dots, n+m-1\}$ find a continuous map $\gamma_i : [i, i+1] \to Y$ such that $\gamma_i(i) = y_i$ and $\gamma_i(i+1) = y_{i+1}$. The maps γ_i compose a single continuous map $\gamma : [1, n+m] \to Y$ such that $\gamma|_{[i,i+1]} = \gamma_i$ for all $i \in \{1, \dots, n+m-1\}$. We define a map $h : X \to Y$,

$$h(x) = \begin{cases} \gamma(\tilde{\psi}(x)), & x \in X \setminus (U \cup V), \\ f(x), & x \in U, \\ g(x), & x \in V. \end{cases}$$

Notice that h is continuous, $h|_A = f|_A$ and $h|_B = g|_B$.

(3). *Necessity.* Fix $y^* \in Y$. Consider the space $X = Y \times [0,1]$, its disjoint functionally closed subsets $A = Y \times \{0\}$ and $B = Y \times \{1\}$, and continuous maps $f, g : X \to Y$ such that $f(y,t) = y$ and $g(y,t) = y^*$ for all $y \in Y$ and $t \in [0,1]$. Since Y is adhesive for X, there exists a continuous map $\lambda : X \to Y$ such that $\lambda|_A = f|_A$ and $\lambda|_B = g|_B$. This implies that Y is contractible.

Sufficiency. Assume that Y is contractible and let $\lambda : Y \times [0,1] \to Y$ be a continuous map such that $\lambda(y,0) = y$ and $\lambda(y,1) = y_0$ for all $y \in Y$. Consider a topological space X, its disjoint functionally closed subsets A and B, and continuous maps $f, g : X \to Y$. Let $\varphi : X \to [0,1]$ be a continuous function such that $A = \varphi^{-1}(0)$ and $B = \varphi^{-1}(1)$. Then a map $h : X \to Y$,

$$h(x) = \begin{cases} \lambda(f(x), 2\varphi(x)), & \varphi(x) \in [0, \frac{1}{2}], \\ \lambda(g(x), 2 - 2\varphi(x)), & \varphi(x) \in (\frac{1}{2}, 1], \end{cases}$$

is continuous and satisfies conditions from Definition 2.1. □

The following example indicates the essentiality of path-connectedness in Theorem 2.7 (2).

Example 2.8. *There exists a connected set $Y \subseteq \mathbb{R}^2$ such that for any two distinct continuous functions $f, g : [0,1] \to Y$ and any two disjoint closed sets A and B in $[0,1]$ there is no continuous function $h : [0,1] \to Y$ with $h|_A = f|_A$ and $h|_B = g|_B$.*

Proof. Let $\mathbb{Q} = \{r_n : n \in \mathbb{N}\}$ be the set of all rational numbers. For every $n \in \mathbb{N}$ we consider the function $\varphi_n : \mathbb{R} \to \mathbb{R}$,

$$\varphi_n(x) = \begin{cases} \sin \frac{1}{x - r_n}, & x \neq r_n, \\ 0, & x = r_n. \end{cases}$$

Define the function $\varphi : \mathbb{R} \to \mathbb{R}$,

$$\varphi(x) = \sum_{n=1}^{\infty} \frac{1}{2^n} \varphi_n(x).$$

Let

$$Y = \{(x, y) \in \mathbb{R}^2 : y = \varphi(x)\}.$$

Observe that for every n the function $\psi_n(x) = \sum\limits_{k=1}^{n} \frac{1}{2^k} \varphi_k(x)$ is a Baire-one Darboux function. Since the sequence $(\psi_n)_{n=1}^{\infty}$ is uniformly convergent to φ on $\mathbb{R}$, φ is a Baire-one Darboux function [3, Theorem 3.4]. Consequently, the graph Y of φ is connected according to [3, Theorem 1.1]. Notice that the space Y is punctiform (i.e., Y does not contain any continuum of the cardinality larger than one), since φ is discontinuous on everywhere dense set $\mathbb{Q}$ (see [10]). Then each continuous function between $\mathbb{R}$ and Y is constant. The statement of the example follows immediately. □

Example 2.9. a) Let $X = [0,1]$, $C \subseteq [0,1]$ be the Cantor set and $Y = \Delta(C) \subseteq \mathbb{R}^2$ be the cone over C, i.e., the collection of all segments which connect the point $v = (\frac{1}{2}, 1)$ with points $(x, 0)$ for all $x \in C$. Then, being contractible, $Y \in \mathrm{AAd}$. We show that Y is not an extensor for X. Indeed, denote by $((a_n, b_n))_{n=1}^{\infty}$ the sequence of contiguous intervals to the Cantor set and consider the identity embedding $f : C \to C \times \{0\}$. Assume that there exists a continuous extension $g : [0,1] \to Y$ of the function f. Then $g([a_n, b_n]) \supseteq [a_n, v] \cup [b_n, v]$ for every $n \in \mathbb{N}$, which implies the equality $g([0,1]) = Y$. Therefore, we obtain a contradiction, since Y is not locally path-connected.

b) Let $X = Y = S^1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$. By Theorem 2.7 the space S^1 is adhesive for itself. On the other hand, the continuous maps $f, g : S^1 \to S^1$ defined by the equalities $f(x, y) = (x, y)$ and $g(x, y) = (1, 0)$, are not homotopic.

3. Stable Baire classes and their characterization

We omit the proof of the following fact, since it is completely similar to the proof of Theorem 2 from [9, p. 357].

Lemma 3.1. *If A is a functionally ambiguous set of the class $\alpha > 1$ in a topological space X, then there exists a sequence $(A_n)_{n=1}^{\infty}$ of functionally ambiguous sets of classes $< \alpha$ such that*

$$A = \lim_{n\to\infty} A_n = \bigcup_{n=1}^{\infty} \bigcap_{k=0}^{\infty} A_{n+k} = \bigcap_{n=1}^{\infty} \bigcup_{k=0}^{\infty} A_{n+k}. \tag{3.1}$$

Moreover, if $\alpha = \beta + 1$ for a limit ordinal β, then all the sets A_n can be taken from classes $< \beta$.

Theorem 3.2. *Let X be a topological space, Y be a topological space with the functionally closed diagonal $\Delta = \{(y, y) : y \in Y\}$, $\alpha \in [1, \omega_1)$ and let $\beta = \alpha$, if $\alpha < \omega_0$, and $\beta = \alpha + 1$, if $\alpha \geq \omega_0$. For a map $f : X \to Y$ we consider the following conditions:*

1. *$f \in \mathrm{B}^{\mathrm{st}}_{\alpha}(X, Y)$;*
2. *there exist an increasing sequence $(X_n)_{n=1}^{\infty}$ of sets of functionally multiplicative classes $< \beta$ and a sequence $(f_n)_{n=1}^{\infty}$ of maps $f_n \in \mathrm{B}^{\mathrm{st}}_{<\alpha}(X, Y)$ such that $X = \bigcup_{n=1}^{\infty} X_n$ $f_n|_{X_n} = f|_{X_n}$ for all $n \in \mathbb{N}$;*
3. *there exist a partition $(X_n : n \in \mathbb{N})$ of X by functionally ambiguous sets of the class β and a sequence of continuous maps $f_n : X \to Y$ such that $f_n|_{X_n} = f|_{X_n}$ for all $n \in \mathbb{N}$.*

Then (1) $\Leftrightarrow$ (2) $\Rightarrow$ (3). If one of the following conditions hold

(a) *Y is adhesive for X, or*
(b) *Y is path-connected σ-adhesive for X,*

then (3) $\Rightarrow$ (2).

Proof. **(1)$\Rightarrow$(2).** Assume that the diagonal Δ is functionally closed in Y^2. Let $(f_n)_{n=1}^{\infty}$ be a sequence of maps $f_n \in \mathrm{B}^{\mathrm{st}}_{<\alpha}(X, Y)$ which is stably convergent to f on X. For $k, n \in \mathbb{N}$ we put

$$X_{k,n} = \{x \in X : f_k(x) = f_n(x)\} \quad \text{and} \quad X_n = \bigcap_{k=n}^{\infty} X_{k,n}.$$

Clearly, $X_n \subseteq X_{n+1}$, $X = \bigcup\limits_{n=1}^{\infty} X_n$ and $f_n|_{X_n} = f|_{X_n}$ for every $n \in \mathbb{N}$.

For all $x \in X$ and $k, n \in \mathbb{N}$ we put $h_{k,n}(x) = (f_k(x), f_n(x))$. In the case $\alpha = \gamma + 1 < \omega_0$ we can assume that $f_n \in \mathrm{B}^{\mathrm{st}}_{\gamma}(X, Y)$ for all $n \in \mathbb{N}$. Then the equality

$$X_{k,n} = h^{-1}_{k,n}(\Delta)$$

implies that $X_{k,n} \in \mathcal{M}_{\gamma}$ and $X_n \in \mathcal{M}_{\gamma}$. Suppose $\alpha \geq \omega_0$. If $\alpha = \omega_0$, then each map f_n can be taken from the class $\mathrm{B}^{\mathrm{st}}_n(X, Y)$. Then $X_{k,n} \in \mathcal{M}_k$ for all $k \geq n$, which implies that $X_n \in \mathcal{M}_{\omega_0}$. Now let $\alpha > \omega_0$. We can assume that $f_n \in \mathrm{B}^{\mathrm{st}}_{\alpha_n}(X, Y)$, where $\omega_0 \leq \alpha_n < \alpha_n + 1 \leq \alpha$ for all $n \in \mathbb{N}$. Then $X_{k,n} \in \mathcal{M}_{\max\{\alpha_n, \alpha_k\}+1} \subseteq \mathcal{M}_{\alpha}$ and $X_n \in \mathcal{M}_{\alpha}$.

(2)⇒(1). Since the sequence $(X_n)_{n=1}^{\infty}$ is increasing, $(f_n)_{n=1}^{\infty}$ is convergent stably to f on X.

(2)⇒(3). We will argue by the induction. For $\alpha = 1$ we take a sequence $(F_n)_{n=1}^{\infty}$ of functionally closed sets and a sequence $(f_n)_{n=1}^{\infty}$ of continuous maps $f_n : X \to Y$ such that $f_n|_{F_n} = f|_{F_n}$ and $X = \bigcup\limits_{n=1}^{\infty} F_n$. We set $X_1 = F_1$ and $X_n = F_n \setminus (F_1 \cup \cdots \cup F_{n-1})$ for $n \geq 2$. Then the family $(X_n : n \in \mathbb{N})$ is the required partition of the space X.

Assume that the implication (2)⇒(3) holds for all $\gamma < \alpha$ for some $\alpha \in [1, \omega_0)$. Let $(A_n)_{n=1}^{\infty}$ be an increasing sequence of sets of the $(\alpha-1)$'th functionally multiplicative class and let $(g_n)_{n=1}^{\infty}$ be a sequence of maps from the class $\mathrm{B}^{\mathrm{st}}_{\alpha-1}(X, Y)$ such that

$$X = \bigcup_{n=1}^{\infty} A_n \quad \text{and} \quad g_n|_{A_n} = f|_{A_n} \quad \text{for every } n \in \mathbb{N} \tag{3.2}$$

By the inductive assumption and by equivalence (1) ⇔ (2), for every n there exist a sequence $(B_{nm})_{m=1}^{\infty}$ of mutually disjoint functionally ambiguous sets of the class $\alpha - 1$ and a sequence $(h_{nm})_{m=1}^{\infty}$ of continuous maps $h_{nm} : X \to Y$ such that

$$h_{nm}|_{B_{nm}} = g_n|_{B_{nm}} \quad \text{for all} \quad m \in \mathbb{N}. \tag{3.3}$$

For all $n, m \in \mathbb{N}$ we set

$$X_{nm} = (A_n \setminus \bigcup_{k=0}^{n-1} A_k) \cap B_{nm}, \tag{3.4}$$

where $A_0 = \emptyset$. Then the partition $(X_{nm} : n, m \in \mathbb{N})$ of the space X is the required one.

We show that (2)⇒(3) for all $\alpha \in [\omega_0, \omega_1)$ under the assumption that Y has the functionally closed diagonal. Again we will argue by the transfinite induction. Let $\alpha = \omega_0$, $(A_n)_{n=1}^{\infty}$ be an increasing sequence of sets of the ω_0'th functionally multiplicative class and $(g_n)_{n=1}^{\infty}$ be a sequence of maps $g_n \in \mathrm{B}^{\mathrm{st}}_n(X, Y)$ such that (3.2) holds. By implication (1)⇒(2) proved above for g_n and by implication (2)⇒(3) proved above for finite ordinals, we obtain that for every $n \in \mathbb{N}$ there exist a partition $(B_{nm})_{m=1}^{\infty}$ of the space X by functionally ambiguous sets of the class n and a sequence $(h_{nm})_{m=1}^{\infty}$ of continuous maps such that (3.3) is valid. It remains to apply (3.4). Further, the inductive step is proved similarly to the case of finite ordinals.

Now we prove that **(3)⇒(2)** in the case $\alpha = 1$. Assume that condition (a) holds. For every $n \in \mathbb{N}$ we take an increasing sequence $(F_{nm})_{m=1}^{\infty}$ of functionally closed

sets in X such that $X_n = \bigcup_{m=1}^{\infty} F_{nm}$ and set $\tilde{X}_n = \bigcup_{m=1}^{n} F_{mn}$. Then $(\tilde{X}_n)_{n=1}^{\infty}$ is an increasing sequence of functionally closed sets which covers the space X. Since Y is adhesive for X, for every $n \in \mathbb{N}$ there exists a continuous map $\tilde{f}_n : X \to Y$ such that $\tilde{f}_n|_{F_{mn}} = f_m|_{F_{mn}}$ for all $m \in \{1, \dots, n\}$. Clearly, $\tilde{f}_n|_{\tilde{X}_n} = f|_{\tilde{X}_n}$ for all $n \in \mathbb{N}$.

Now suppose that condition **(b)** holds. Let Y be a path-connected σ-adhesive for X and $(Y_n : n \in \mathbb{N})$ be a cover of the space Y by functionally closed subspaces Y_n such that $(Y, Y_n) \in \mathrm{Ad}^*(X)$ for every n.

We prove that the preimage of each functionally closed subset of Y under the mapping f is functionally ambiguous of the class β in X. Indeed, take a functionally closed set $B \subseteq Y$. Then $f^{-1}(B) = \bigcup_{n=1}^{\infty} (f_n^{-1}(B) \cap X_n)$. Since $f_n : X \to Y$ is continuous, $f_n^{-1}(B)$ is functionally closed in X. Therefore, $f^{-1}(B)$ belongs to the β'th additive class in X. Moreover, $X \setminus f^{-1}(B) = \bigcup_{n=1}^{\infty} (f_n^{-1}(Y \setminus B) \cap X_n)$. Since $f_n^{-1}(Y \setminus B)$ is functionally open in X, we have that $X \setminus f^{-1}(B)$ belongs to the functionally additive class β in X. Thus, $f^{-1}(B)$ is functionally ambiguous of the β'th class in X.

For every $k, n \in \mathbb{N}$ we put $X_{k,n} = f^{-1}(Y_k) \cap X_n$. Let us remove from the sequence $(X_{k,n})_{k,n=1}^{\infty}$ empty sets and let $(Z_n)_{n=1}^{\infty}$ be an enumeration of the double sequence. Denote $\tilde{X}_1 = Z_1$ and $\tilde{X}_n = Z_n \setminus \bigcup_{k<n} Z_k$ for $n > 1$. Notice that $(\tilde{X}_n : n \in \mathbb{N})$ is a partition of X by functionally ambiguous sets of the class β in X. For every $k \in \mathbb{N}$ we set $N_k = \{n \in \mathbb{N} : f(\tilde{X}_n) \subseteq Y_k\}$ and put $\tilde{Y}_i = Y_k$ for all $i \in N_k$. Hence, we obtain a partition $(\tilde{X}_n : n \in \mathbb{N})$ of X by functionally ambiguous sets of the β'th class in X such that $f(\tilde{X}_n) \subseteq \tilde{Y}_n$ for every n and it is evident that $(Y_n : n \in \mathbb{N})$ has the same properties as $(Y_n : n \in \mathbb{N})$.

For every $n \in \mathbb{N}$ we take an increasing sequence $(F_{nm})_{m=1}^{\infty}$ of functionally closed sets in X such that $\tilde{X}_n = \bigcup_{m=1}^{\infty} F_{nm}$ and denote $C_n = \bigcup_{m=1}^{n} F_{mn}$. We fix $n \in \mathbb{N}$ and show that the restriction $f|_{C_n} : C_n \to Y$ has a continuous extension $g : X \to Y$.

Since the sets $F_{1n}, \dots, F_{nn}$ are disjoint and functionally closed, we may choose open sets $U_1, \dots, U_n$ and a continuous function $\varphi : X \to [1, n]$ such that $F_{mn} \subseteq U_m$ and $\overline{U}_m \subseteq \varphi^{-1}(m)$ for every $m \in \{1, \dots, n\}$. Further, in each $\tilde{Y}_m$ we take a point y_m^* from Definition 2.3. Since $(Y, \tilde{Y}_m) \in \mathrm{Ad}^*(X)$, there exists a continuous map $g_m : X \to Y$ such that $g_m|_{F_{mn}} = f|_{F_{mn}}$ and $g_m|_{X \setminus U_m} = y_m^*$. It implies from the path-connectedness of Y that there exists a continuous map $\gamma : [1, +\infty) \to Y$ such that $\gamma(m) = y_m^*$ for every $m \in \mathbb{N}$. For all $x \in X$ we set

$$g(x) = \begin{cases} g_m(x), & \text{if } x \in U_m \text{ for some } m \in \{1, \dots, n\}, \\ \gamma(\varphi(x)), & \text{otherwise.} \end{cases}$$

Then the map $g : X \to Y$ is continuous and $g|_{C_n} = f|_{C_n}$. Hence, condition (2) holds.

Now we suppose that under conditions (a) or (b) the implication (3)⇒(1) is valid for all ordinals $\gamma \in [1, \alpha)$ for some $\alpha \in (1, \omega_1)$ and prove it for α. Consider the case $\alpha = \gamma + 1 < \omega_0$. By Lemma 3.1 for every m there exists a sequence $(A_{mn})_{n=1}^{\infty}$ of functionally ambiguous sets of the class γ such that $X_m = \operatorname{Lim}_{n\to\infty} A_{mn}$. For all

$m, n \in \mathbb{N}$ we set

$$B_{mn} = A_{mn} \setminus \bigcup_{k<m} A_{kn}. \tag{3.5}$$

Then each set B_{mn} is functionally ambiguous of the class γ.

For every $n \in \mathbb{N}$ we set

$$g_n(x) = \begin{cases} f_m(x), & \text{if } x \in B_{mn} \text{ for } m < n, \\ f_n(x), & \text{otherwise} \end{cases}$$

and, applying the inductive assumption, we get $g_n \in \mathrm{B}^{\mathrm{st}}_{\gamma}(X, Y)$.

It remains to prove that $(g_n)_{n=1}^{\infty}$ is stably convergent to f on X. Fix $x \in X$ and choose a number m such that $x \in X_m$ and $x \notin X_k$ for all $k \neq m$. Equality (3.1) implies that there are numbers $N_1, \dots, N_m$ such that

$$x \notin \bigcup_{n \geq N_k} A_{kn} \text{ if } k < m \text{ and } x \in \bigcap_{n \geq N_m} A_{mn}.$$

Hence, for all $n \geq n_0 = \max\{N_1, \dots, N_m\}$ the inclusion $x \in B_{mn} \cap X_m$ holds. Therefore, $g_n(x) = f_m(x) = f(x)$ for all $n \geq n_0$.

In the case $\alpha \geq \omega_0$ we observe that each set X_n is functionally ambiguous of the class $\alpha + 1$ and the sets A_{mn} (together with the sets B_{mn}) are functionally ambiguous either of the class α, or of classes $< \alpha$ in the case of limit α. Then by the inductive assumption we have $g_n \in \mathrm{B}^{\mathrm{st}}_{\gamma}(X, Y)$, if $\alpha = \gamma + 1 > \omega_0$, and $g_n \in \mathrm{B}^{\mathrm{st}}_{<\alpha}(X, Y)$, if α is limit. In any case, $f \in \mathrm{B}^{\mathrm{st}}_{\alpha}(X, Y)$. □

Let us observe that in the proof of implication (2)⇒(1) we do not use the fact that Y has the functionally closed diagonal. The following example show that this property is essential for implication (1)⇒(2).

Example 3.3. *Let D be an uncountable discrete space and $X = Y = D \sqcup \{a\}$ be the Alexandroff compactification of D. Then there exists $f \in \mathrm{B}^{\mathrm{st}}_1(X, Y)$ such that for every functionally measurable subset $A \subseteq X$ with $a \in A$ the restriction $f|_A$ is discontinuous at a.*

Proof. Let $D = \bigsqcup_{n=1}^{\infty} D_n$ such that $|D| = |D_n|$ for every $n \in \mathbb{N}$. We choose a sequence of bijections $\varphi_n : D_n \to D$ and consider the function $f : X \to Y$,

$$f(x) = \begin{cases} a, & x = a, \\ \varphi_n(x), & n \in \mathbb{N} \text{ and } x \in D_n. \end{cases}$$

Note that $f \in \mathrm{B}^{\mathrm{st}}_1(X, Y)$, because f is the stable limit of the sequence of continuous functions $f_n : X \to Y$,

$$f_n(x) = \begin{cases} a, & x \in \{a\} \cup (\bigcup_{k=n+1}^{\infty} D_k), \\ \varphi_k(x), & k \leq n \text{ and } x \in D_k. \end{cases}$$

Fix a functionally measurable subset $A \ni a$ of X. Since $|D \setminus B| \leq \aleph_0$ for every functionally open or functionally closed subset $B \ni a$ of X, we have $|D \setminus A| \leq \aleph_0$. Thus every set $B_n = D_n \setminus A$ is at most countable. Therefore the set $C = \bigcup_{n=1}^{\infty} \varphi_n(B_n)$ is at most countable too. We choose $d \in D \setminus C$ and consider the neighborhood $V = Y \setminus \{d\}$ of a in Y. Since

$$A \setminus \{x \in A : f(x) \in V\} = \{\varphi_n^{-1}(d) : n \in \mathbb{N}\},$$

the restriction $f|_A$ is discontinuous at a. □

We show in the following example that the properties (a) and (b) in Theorem 3.2 are essential.

Example 3.4. a) Let $X = [0,1]^2$ and $Y \subseteq X$ be the Sierpiński carpet. Notice that Y is a Peano continuum. Fix any $x^* \in Y$ and consider the map $f : X \to Y$ such that $f(x) = x$ for $x \in Y$ and $f(x) = x^*$ for $x \in X \setminus Y$. We put $f_1(x) = x$ and $f_2(x) = x^*$ for all $x \in X$. Notice that $X_1 = Y$ and $X_2 = X \setminus Y$ are ambiguous subsets of the first class in X, $X_1 \cup X_2 = X$ and $f|_{X_i} = f_i|_{X_i}$ for $i = 1, 2$. Therefore, condition (3) of Theorem 3.2 holds.

Assume that $f \in \mathrm{B}_1^{\mathrm{st}}(X, Y)$. Take a sequence $(g_n)_{n=1}^{\infty}$ of continuous functions $g_n : X \to Y$ and a closed cover $(\tilde{X}_n : n \in \mathbb{N})$ of X such that $g_n|_{\tilde{X}_n} = f|_{\tilde{X}_n}$ and $\tilde{X}_n \subseteq \tilde{X}_{n+1}$ for every n. By the Baire category theorem, there exists $k \in \mathbb{N}$ such that the set $\tilde{X}_k \cap Y$ has the nonempty interior in Y. Then there exists an open square L in Y such that $\partial L \subseteq \tilde{X}_k \cap Y$. Since $g_k|_{\partial L}(x) = x$ for all $x \in \partial L$ and $g_k(\mathrm{int} L) \subseteq X \setminus \mathrm{int} L$, we have that ∂L is a retract of $\overline{L}$, which is impossible. Hence, condition (2) of Theorem 3.2 does not hold. Consequently, Y fails to be adhesive for X.

b) Let $Y = \{0, 1\}$ and $f : [0,1] \to Y$ be the characteristic function of the set $\{0\}$. Clearly, Y is a disconnected σ-adhesive space for $[0,1]$. Moreover, the partition $(\{0\}, (0,1])$ of $[0,1]$ and the functions $f_1 \equiv 1$ and $f_2 \equiv 0$ satisfy condition (3) of Theorem 3.2, but f is not of the first stable Baire class, since each continuous function between $[0,1]$ and $\{0,1\}$ is constant.

Clearly, if $\alpha = 1$, then condition (3) of Theorem 3.2 implies that f is piecewise continuous. It was proved in [1, Theorem 6.3] that every piecewise continuous map belongs to the class $\mathrm{B}_1^{\mathrm{st}}(X, Y)$ when X is a normal space and Y is a path-connected space such that $Y \in \sigma\mathrm{AE}(X)$. The following example shows that a piecewise continuous map need not be of the first stable Baire class even if $X = \mathbb{R}^2$ and Y is a contractible subspace of $\mathbb{R}^2$.

Example 3.5. *Let $Y = \Delta(C)$ be the cone over the Cantor set $C \subseteq [0,1]$ defined in Example 2.9, $y^* \in Y$ be a point and $f : [0,1]^2 \to Y$ be a map such that*

$$f(x) = \begin{cases} x, & x \in Y, \\ y^*, & \textit{otherwise.} \end{cases}$$

Then f is piecewise continuous and $f \notin \mathrm{B}_1^{\mathrm{st}}([0,1]^2, Y)$.

Proof. Since $f|_Y$ and $f|_{[0,1]^2 \setminus Y}$ is continuous and Y is closed in $[0,1]^2$, f is piecewise continuous.

Assume that there exist a sequence $(f_n)_{n=1}^{\infty}$ of continuous functions $f_n : [0,1]^2 \to Y$ and a closed cover $(X_n : n \in \mathbb{N})$ of the square $[0,1]^2$ such that $f_n|_{X_n} = f|_{X_n}$ for every n. By the Baire category theorem, there exists $k \in \mathbb{N}$ such that the set $X_k \cap Y$ has the nonempty interior in Y. Let F be a closed square in $[0,1]^2$ such that $F \cap Y \subseteq X_k \cap Y$ and the interior of $F \cap Y$ in Y is nonempty. Consider the set $B = f_k(F)$. Since B is a continuous image of F, it should be locally connected. On the other hand, B is not a locally connected set, since B is a closed subspace of Y with nonempty interior. □

4. Monotone functions and their stable Baire measurability

We will establish in this section that

$$\mathcal{M} \cap \mathrm{B}_1^{\mathrm{st}}(\mathbb{R},\mathbb{R}) = \mathcal{M} \cap \mathrm{B}_2^{\mathrm{st}}(\mathbb{R},\mathbb{R}) = \cdots = \mathcal{M} \cap \mathrm{B}_\alpha^{\mathrm{st}}(\mathbb{R},\mathbb{R}) = \ldots,$$

where $\mathcal{M}$ is the class of all monotone functions.

The following fact immediately follows from definitions and we omit its proof.

Lemma 4.1. *Let $X \subseteq \mathbb{R}$, $f : X \to \mathbb{R}$ be a monotone function and $g : X \to \mathbb{R}$ be a continuous function such that $f|_D = g|_D$ for some dense set $D \subseteq X$. Then $f = g$ on X.*

A map $f : X \to Y$ between topological spaces X and Y is said to be *weakly discontinuous*, if for any subset $A \subseteq X$ the discontinuity points set of the restriction $f|_A$ is nowhere dense in A. It is easy to see that a map f is weakly discontinuous if and only if the discontinuity points set of the restriction $f|_F$ to any closed set $F \subseteq X$ is nowhere dense in F.

Theorem 4.2. *For a monotone function $f : \mathbb{R} \to \mathbb{R}$ the following conditions are equivalent:*

(1) *f is weakly discontinuous;*
(2) *$f \in \mathrm{B}_1^{\mathrm{st}}(\mathbb{R},\mathbb{R})$;*
(3) *$f \in \bigcup\limits_{\alpha<\omega_1} \mathrm{B}_\alpha^{\mathrm{st}}(\mathbb{R},\mathbb{R})$.*

Proof. The equivalence of (1) and (2) was established in [2] (see also [1] for a more general case).

The implication (2)⇒(3) is evident.

We prove that (3)⇒(1). Let $f \in \mathrm{B}_\alpha^{\mathrm{st}}(\mathbb{R},\mathbb{R})$ for some $\alpha \in [0,\omega_1)$. By Theorem 3.2 there exist a sequence of continuous functions $f_n : \mathbb{R} \to \mathbb{R}$ and a partition $(X_n : n \in \mathbb{N})$ of the real line such that $f_n|_{X_n} = f|_{X_n}$ for every $n \in \mathbb{N}$. Consider a nonempty closed set $F \subseteq \mathbb{R}$. For every n we denote $G_n = \mathrm{int}_F \overline{X_n \cap F}$. Since F is a Baire space, the union $G = \bigcup\limits_{n=1}^{\infty} G_n$ is dense in F. The equality $f|_{X_n \cap F} = f_n|_{X_n \cap F}$ and Lemma 4.1 imply that $f|_{G_n} = f_n|_{G_n}$ for every n. Since every function f_n is continuous and the set G_n is open in F, we have $G_n \subseteq C(f|_F)$. Hence, $G \subseteq C(f|_F)$, which implies that f is weakly discontinuous. □

As a corollary of Theorem 4.2 we obtain the following result.

Proposition 4.3. *There exists a function $f \in \mathrm{B}_1(\mathbb{R},\mathbb{R}) \setminus \bigcup\limits_{\alpha<\omega_1} \mathrm{B}_\alpha^{\mathrm{st}}(\mathbb{R},\mathbb{R})$.*

Proof. We consider the increasing function $f : \mathbb{R} \to \mathbb{R}$,

$$f(x) = \sum_{r_n \le x} \frac{1}{2^n},$$

where $\mathbb{Q} = \{r_n : n \in \mathbb{N}\}$. Since f is monotone, $f \in \mathrm{B}_1(\mathbb{R},\mathbb{R})$. But $D(f) = \mathbb{Q}$. Therefore, f is not weakly discontinuous. Hence, $f \notin \bigcup\limits_{\alpha<\omega_1} \mathrm{B}_\alpha^{\mathrm{st}}(\mathbb{R},\mathbb{R})$ by Theorem 4.2. □

References

[1] T. Banakh, B. Bokalo *On scatteredly continuous maps between topological spaces*, Topology Appl. **157** (1) (2010), 108-122.

[2] T. Banakh, S. Kutsak, O. Maslyuchenko, V. Maslyuchenko, *Direct and inverse problems of the Baire classifications of integrals dependent on a parameter*, Ukr. Math. J. **56** (11) (2004), 1443–1457.

[3] A. Bruckner, *Differentiation of Real Functions* [2nd ed.], Providence, RI: American Mathematical Society, 1994, 195 p.

[4] Á. Császár, M. Laczkovich, *Discrete and equal convergence*, Studia Sci. Math. Hungar. **10** (1975), 463–472.

[5] Á. Császár, M. Laczkovich, *Some remarks on discrete Baire classes*, Acta Math. Acad. Sci. Hungar. **33** (1979), 51–70.

[6] F. Chaatit, H. Rosenthal, *On differences of semi-continuous functions*, Quaest. Math. **23** (3) (2000) 295–311.

[7] R. Haydon, E. Odell, H. Rosenthal, *On certain classes of Baire-1 functions with applications to Banach space theory*, in: Functional Analysis, Austin, TX, 1987/1989, in: Lecture Notes in Math., vol. 1470, Springer, Berlin, 1991, pp. 1–35.

[8] O. Karlova, V. Mykhaylyuk, *On composition of Baire functions*, arxiv.org/abs/1511.08982.

[9] K. Kuratowski, *Topology. Volume 1*, Academic Press (1966).

[10] C. Kuratowski and W. Sierpinski, *Les fonctions de classe et les ensembles connexes ponctiformes*, Fund. Math. 3 (1922) 303–313.

E-mail address: maslenizza.ua@gmail.com

E-mail address: vmykhaylyuk@ukr.net

Yurii Fedkovych Chernivtsi National University, Ukraine